

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : A61K 31/00	A2	(11) International Publication Number: WO 98/25593 (43) International Publication Date: 18 June 1998 (18.06.98)
(21) International Application Number: PCT/EP97/06930 (22) International Filing Date: 11 December 1997 (11.12.97) (30) Priority Data: RM96A000867 13 December 1996 (13.12.96) IT (71) Applicant (for all designated States except US): CONSIGLIO NAZIONALE DELLE RICERCHE [IT/IT]; Piazzale Aldo Moro, 7, I-00198 Roma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): SANTORO, Maria, Gabriella [IT/IT]; Via Vasto, 8, I-83100 Avellino (IT). ROSSI, Antonio [IT/IT]; Via Orientale, 33, I-66010 Colledara (IT). ELIA, Giuliano [IT/IT]; Via della Stella, 321, I-00036 Palestrina (IT). (74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi, Corso di Porta Vittoria, 9, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: 2-CYCLOPENTEN-1-ONE AND ITS DERIVATIVES AS INHIBITORS OF THE NF- κ B FACTOR (57) Abstract 2-Cyclopenten-1-one and its derivatives comprising the cyclopentenone nucleus as inhibitors of the NF- κ B factor, with anti-inflammatory, anti-proliferative, immuno-suppressive, cytoprotective and antiviral activity, the substituents being selected among the ones which do not affect the NF- κ B inhibitory activity.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MR	Mauritania	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MW	Malawi	UA	Ukraine
BR	Brazil	IS	Iceland	MX	Mexico	UG	Uganda
BY	Belarus	IT	Italy	NE	Niger	US	United States of America
CA	Canada	JP	Japan	NL	Netherlands	UZ	Uzbekistan
CF	Central African Republic	KE	Kenya	NO	Norway	VN	Viet Nam
CG	Congo	KG	Kyrgyzstan	NZ	New Zealand	YU	Yugoslavia
CH	Switzerland	KP	Democratic People's Republic of Korea	PL	Poland	ZW	Zimbabwe
CI	Côte d'Ivoire			PT	Portugal		
CM	Cameroon	KR	Republic of Korea	RO	Romania		
CN	China	KZ	Kazakstan	RU	Russian Federation		
CU	Cuba	LC	Saint Lucia	SD	Sudan		
CZ	Czech Republic	LI	Liechtenstein	SE	Sweden		
DE	Germany	LK	Sri Lanka	SG	Singapore		
DK	Denmark	LR	Liberia				
EE	Estonia						

2-CYCLOPENTEN-1-ONE AND ITS DERIVATIVES AS INHIBITORS OF THE NF- κ B
FACTOR

FIELD OF INVENTION

The present invention relates to 2-cyclopenten-1-one and its
5 derivatives as inhibitors of the transcription factor Nuclear Factor- κ B (NF- κ B). In particular the invention relates to 2-cyclopenten-1-one and its derivatives comprising the cyclopentenone nucleus as inhibitors of the NF- κ B factor with anti-inflammatory, anti-proliferative, immuno-suppressive, cytoprotective and antiviral
10 activity.

BACKGROUND OF THE INVENTION

NF- κ B (Nuclear Factor - κ B) is an eucariotic transcription factor of the *rel* family, which is normally located in the cytoplasm in an inactive complex, whose predominant form is a heterodimer composed of
15 p50 and p65 subunits, bound to inhibitory proteins of the I κ B family, usually I κ B- α (D. Thanos and T. Maniatis, and Cell 80:529-532, 1995).

NF- κ B is activated in response to different stimuli, among which phorbol esters, inflammatory cytokines, UV radiation, bacterial and
20 viral infections. Stimulation triggers the release of NF- κ B from I κ B in consequence of the phosphorylation and the following degradation of the I κ B- α protein (P.A. Baeuerle and T. Henkel, Annu. Rev. Immunol. 12: 141-179, 1994). Once it is activated, NF- κ B translocates to the nucleus where it binds to DNA at specific κ B-sites and induces
25 the transcription of a variety of genes encoding proteins involved in controlling the immune and inflammatory responses, among which a variety of interleukins, the tumor necrosis factor α , the NO

synthase and the cyclo-oxygenase 2 (S. Grimm and P.A. Baeuerle, Biochem. J. 290: 297-308, 1993). Accordingly, NF-kB is considered an early mediator of the immune and inflammatory responses and it is involved in the control of cell proliferation and in the pathogenesis of various human diseases, among which rheumatoid arthritis (H. Beker et al., Clin. Exp. Immunol. 99: 325, 1995), ischemia (A. Salminen et al. Biochem. Biophys. Res. Comm. 212: 939, 1995), arteriosclerosis (A.S. Baldwin. Annals Rev. Immunol. 14: 649, 1996), as well as in the pathogenesis of the acquired immunodeficiency syndrome (AIDS), due to the enhanced human immunodeficiency virus (HIV-1) transcription in the presence of activated NF-kB. The increase of HIV-1 virus RNAs transcription by NF-kB is caused by the presence of kB-sites in the (LTR) (Long Terminal Repeats) sequences of the virus genome (M.J. Lenardo and D. Baltimore, Cell 58: 227-229, 1989).

It is also known that prostaglandins (PGs) are a class of naturally occurring cyclic 20-carbon fatty acids that are synthesized by various types of eukaryotic cells in response to external stimuli and play an important role in a variety of physiological responses. Since their discovery, PGs were shown to act as microenvironmental hormones and intracellular signal mediators and to control a large number of physiological and pathological processes, including cell proliferation and differentiation, the immune response, inflammation, cytoprotection and the febrile response. In particular, type A and J PGs, which possess a cyclopentenonic structure, are strong inhibitors of virus replication ("Stress Proteins: Induction and Function" Schlesinger MJ, Garaci E., Santoro M.G. ed.s, Springer-Verlag, Heidelberg-Berlin, 27-44, 1990). Particularly, it has been recently demonstrated that

cyclopentenonic prostaglandins inhibit HIV-1 virus replication, by blocking the viral RNAs transcription (C. Rozera et al. J. Clin. Invest. 97: 1795, 1996).

It is also known that the Heat Shock Proteins (HSPs), also called
5 stress proteins (Proc. Natl. Acad. Sci. USA 86, 8407-8411, 1989), are a family of polypeptides synthesized by eukaryotic and prokaryotic cells in response to heat shock or other kinds of environmental stresses. The HSPs are encoded by a cellular subgroup of genes, identified as stress genes.

10 The authors have shown that the cyclopentenone prostaglandin PGA inhibits the activation of NF- κ B in human cells by inhibiting the phosphorylation and degradation of the inhibitory I κ B- α protein (A. Rossi, G. Elia and M.G. Santoro, Cold Spring Harbour, New York 1-5 May, 1996, Abstract p. 255).

15 The authors have also recently shown that inhibition of NF- κ B activation is one of the molecular mechanisms used by cyclopentenonic prostaglandins to cause a selective and reversible block of HIV-1 virus RNAs transcription.

SUMMARY OF THE INVENTION

20 It has now been found that 2-cyclopenten-1-one, the structure constituting the center nucleus of PGA, possesses an activity which is analogous to PGA, that is, it is able to inhibit NF- κ B activation, even though it does not contain the corresponding acid function and aliphatic lateral chains. Therefore it is found that the lateral
25 chains, which are present in the PGA with their substituents and double bonds, in particular the acid function, which implies the fatty acid nature of prostaglandins, can be eliminated without substantially

modifying the herein above described specific activity. It is also found that the α,β -unsaturated carbonyl group in the cyclopentenone ring is the key structure necessary for NF- κ B inhibition.

Furthermore it has been found that the inhibition of NF- κ B by the
5 cyclopentenone group is related to the ability to activate the HSF transcription factor (Heat Shock Transcription Factor), which is responsible for the synthesis of HSPs (Heat Shock Proteins).

In view of the fact that NF- κ B inhibition is associated with HSF activation, it is evident that molecules containing the cyclopentenone
10 nucleus, which is active in inhibiting NF- κ B, will be inducers of the HSF factor and therefore they will be inducers of heat shock proteins. It is therefore an object of the present invention the 2-cyclopenten-1-one, and its substituted derivatives comprising the cyclopentenone nucleus, as inhibitors of NF- κ B, the substituents being selected among
15 the ones which do not affect the NF- κ B inhibitory activity.

Another object of the present invention is the 2-cyclopenten-1-one and its pharmacologically acceptable derivatives as inhibitors of NF- κ B.

Another object of the invention is the 2-cyclopenten-1-one and its derivatives as inhibitors of NF- κ B with anti-inflammatory, anti-
20 proliferative, immuno-suppressive, cytoprotective and antiviral activity.

A further object of the invention are pharmaceutical compositions comprising 2-cyclopenten-1-one and/or its pharmaceutically acceptable derivatives to make medicaments with anti-inflammatory, anti-
25 proliferative, immuno-suppressive, cytoprotective and antiviral activity. In particular with antiviral activity against the HIV-1 virus and viruses whose transcription is controlled by NF- κ B,

including herpesviruses.

BRIEF DESCRIPTION OF FIGURES

Fig. 1A shows the dose-dependent inhibition of NF-kB activation by 2-cyclopenten-1-one.

- 5 Fig. 1B shows the activation of the HSF factor (Heat Shock Transcription Factor) by 2-cyclopenten-1-one in association with NF-kB inhibition.

Fig. 1C shows the specificity of the chemical structure which is responsible for NF-kB inhibition.

- 10 Fig. 1D shows the specificity of the chemical structure which is responsible for HSF activation.

DETAILED DESCRIPTION OF THE INVENTION

The 2-cyclopenten-1-one is a known product, which can be synthesized according to the process described in Beilstein (*Daene, Eder*; A. 539

- 15 [1939] 207, 211).

According to the present invention 2-cyclopenten-1-one, preferably in concentrations ranging between 100 and 500 μ M, is able to inhibit NF-kB activation in human cells (Fig. 1A).

- Inhibition tests have been carried out in type T lymphoid human cells (Jurkat cell line), as well as in other human cell lines. NF-kB activation was stimulated with 12-o-tetradecanoyl-phorbol-13-acetate (TPA). 2-Cyclopenten-1-one was also effective in inhibiting NF-kB activation after other types of stimulation, including stimulation by tumor necrosis factor alpha or viral infection, and in different types of human cells (data not shown). It is demonstrated that NF-kB inhibition is associated with the activation of HSF factor (Fig. 1B).
- 20
- 25

It is also proved that the ability to inhibit the NF-kB factor is

specific for 2-cyclopenten-1-one, whereas similar molecules, such as cyclopentanone and cyclopentene, do not inhibit NF-kB (Fig. 1C) and do not activate HSF (Fig. 1D).

Based on these results it is possible to use 2-cyclopenten-1-one, as well as its pharmaceutically acceptable derivatives, as active substances to produce medicaments, in particular medicaments having activity in inhibiting the NF-kB factor, and in particular:

- anti-inflammatory and immunosuppressive medicaments, in view of the role of NF-kB in stimulating the inflammatory and immune responses;
- 10 - cytoprotective medicaments, in view of the role of NF-kB in ischemia and oxidative damages;
- antiproliferative medicaments, in view of the role of NF-kB in cell proliferation;
- antiviral medicaments, in view of the role of NF-kB in activating
- 15 the viral RNAs transcription.

The following examples are reported to illustrate the invention. They should be considered in any case non limiting the scope of the invention itself.

The reagents used in the examples, including 2-cyclopenten-1-one, cyclopentene, cyclopentanone and 12-o-tetradecanoyl-phorbol-13-acetate (TPA), were products of Sigma Aldrich. ^{32}P e ^{35}S were produced by AMERSHAM. Fetal calf serum and cellular culture media were produced by GIBCO.

EXAMPLE 1

25 The effect of the treatment with 2-cyclopenten-1-one on NF-kB activation by TPA has been tested in Jurkat cells by using the procedures described hereinbelow and shown in Fig. 1.

Dose-response effect

The cells were prepared according to the method described in C. Amici et al. (Cancer Research 55, 4452-4457, 1995).

The cells were treated with 2-cyclopenten-1-one at different concentrations (125-100uM) for 1 hour and then were stimulated with TPA (25 ng/ml).

After 3 hours the whole-cell extracts were prepared and subjected to EMSA (Electrophoretic Mobility Shift Assay) as described for NF-kB in U. Zabel et al. (J. Biol. Chem. 266: 242, 1991) and HSF in C. Amici et al. (Cancer Res. 55: 4452, 1995), to determine NF-kB and HSF activation respectively. The positions of the complex NF-kB-DNA (NF-kB) and the non-specific binding (ns) are indicated in Fig. 1A.

The positions of the complex HSF-DNA (HSF), the HSF-DNA constitutive activity (CHBA) and the proteins-DNA non-specific interactions (ns) are indicated in Fig. 1B. The line "control" indicates the non-TPA-stimulated cells as a control of non-activated NF-kB.

As evident, 2-cyclopenten-1-one is able to inhibit NF-kB activation by TPA even at the lower concentration of 125 uM. At the concentration of 500 uM the NF-kB band is absent (Fig. A). In correlation with NF-kB inhibition, in the same samples it is evident the activation of HSF starting from the concentration of 125 uM (Fig. 1B).

Specificity of the inhibitory effect (Fig. 1C)

The cells were treated for 1 hour with the same concentration (500 uM) of: 2-cyclopenten-1-one (line 3), cyclopentanone (line 4) or cyclopentene (line 5), and then were stimulated with TPA (25 ng/ml). C represents the non-TPA-stimulated control. After 3 hours the whole-cell extracts were prepared and subjected to EMSA to verify the

activation of NF- κ B (Fig. 1C) and of HSF (Fig. 1D) respectively.

As evident, (i) TPA activates NF- κ B (line 2); (ii) 2-cyclopenten-1-one inhibits TPA-induced NF- κ B activation (line 3); cyclopentanone (line 4) and cyclopentene (line 5) do not inhibit NF- κ B activation.

5 In addition, as shown in Fig. 1D, in the same samples inhibition of NF- κ B, shown in Fig. 1C, is associated with activation of HSF. These results clearly show that the α,β -unsaturated carbonyl group is the key structure triggering HSF activation and its presence is necessary to inhibit NF- κ B activation.

CLAIMS

- 1 1. 2-cyclopente-1-one and its substituted derivatives comprising the
2 cyclopentenonic nucleus as inhibitors of NF-kB factor, the
3 substituents being selected among the ones which do not affect the NF-
4 kB inhibitory activity.
- 1 2. 2-cyclopenten-1-one and its pharmaceutically acceptable derivatives
2 as NF-kB inhibitors with anti-inflammatory, anti-proliferative,
3 immuno-suppressive, cytoprotective and antiviral activity.
- 1 3. 2-cyclopenten-1-one and its pharmaceutically acceptable derivatives
2 to make pharmaceutic compositions with anti-inflammatory, anti-
3 proliferative, immuno-suppressive, cytoprotective and antiviral
4 activity.
- 1 4. 2-cyclopenten-1-one and its pharmaceutically acceptable derivatives
2 as inducers of the HSF factor.
- 1 5. Pharmaceutical compositions comprising 2-cyclopenten-1-one and its
2 pharmaceutically acceptable derivatives to make medicaments with anti-
3 inflammatory, anti-proliferative, immuno-suppressive, cytoprotective
4 and antiviral activity.
- 1 6. Pharmaceutical compositions comprising 2-cyclopenten-1-one and its
2 pharmaceutically acceptable derivatives to make medicaments with
3 antiviral activity against HIV-1 virus and other viruses whose
4 transcription is regulated by the NF-kB factor, including
5 herpesviruses.
- 1 7. Pharmaceutical compositions comprising 2-cyclopenten-1-one and its
2 pharmaceutically acceptable derivatives as inducers of the HSF factor.




FIGURE 1

